

Go to your GP or your nearest Emergency Department immediately if:

- · The wound appears to have re-opened.
- The area looks to be infected (signs include an increase in redness, hot to touch (avoid touching too often with dirty hands), bad smell – pus.
- · You are becoming unwell with a fever.
- You are developing a swelling that affects the ability to eat, drink or swallow.

Information reproduced with permission from Birmingham Women's and Children's NHS Foundation Trust

Further sources of information

NHS Choices: www.nhs.uk/conditions
Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222 Newark Hospital: 01636 685692 Email: sfh-tr.PET@nhs.net

If you would like this information in an alternative format, for example large print or easy read, or if you need help with communicating with us, for example because you use British Sign Language, please let us know. You can call the Patient Experience Team on 01623 672222 or email sfh-tr.PET@nhs.net.

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you. External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them. If you require a full list of references (if relevant) for this leaflet, please email sfh-tr.patientinformation@nhs.net_or telephone 01623 622515, extension 6927.

To be completed by the Communications office Leaflet code: PII 202401-01-FW

Created: January 2024/ Review Date: January 2026

INFORMATION FOR PATIENTS

Facial wounds

Emergency Department

King's Mill Hospital

Mansfield Road Sutton in Ashfield Notts NG17 4.II

Telephone: 01623 622515, extension 2789

Newark Hospital

Boundary Road

Newark

Notts NG24 4DE

Telephone: 01636 681681

Healthier Communities, Outstanding Care A facial wound is when there is an injury or break of facial skin. This can include a cut (laceration), scratches (abrasions), grazes or loss of skin or tissue. This can occur following trauma/assault/ bites.

The deeper layers of the skin are exposed and sometimes become dirty because they have touched things like soil, gravel or even animal saliva.

Treatment

Depending on whether the wound is dirty or gaping open the treatment may consist of:

- 1) Cleaning
- 2) Dressing
- 3) Gluing
- 4) Steri-strips
- Stitching (suturing) either under local anaesthetic, under general anaesthetic, using non-dissolvable stitches or using dissolvable stitches.

Care following the closure of a facial wound

If local anaesthetic was used, the area will be numb for a few hours after treatment so be careful not to pull on the stitches. Keep the area dry and clean for a few days after treatment unless advised to apply ointment

If a dressing has been placed on top of the wound, try to keep this clean and dry. The dressing can be replaced or removed if it is visibly dirty.

General advice:

- Dissolvable stitches usually take two to three weeks to breakdown on their own.
- If non-dissolvable stitches were used, these will need to be removed by your GP in about five to seven days.

- Avoid playing or pulling on the stitches.
- If glue was used to close the wound, this will take a few days to dissolve on its own.
- Steri-strips will peel off on their own a few days after they were placed.
- Don't worry if the area is swollen and puffy after the injury for a few days.
- If you are in pain, take painkillers you usually take for a headache or cold. The area might turn grey/white a few days after the injury. This is normal and is and is often a part of the healing process.

Bite injuries:

- Take antibiotics as directed.
- Keep the wound clean and dry.
- · Apply ointment as directed.